

On Calculation of $2 + \varepsilon$ RG Functions in the Gross Neveu Model from Large N Expansions of Critical Exponents.

Kivel N.A. , Stepanenko A.S.*

*Theoretical Department, St.-Petersburg Nuclear Physics Institute,
Gatchina, St.-Petersburg, 188350, Russian Federation*

Vasil'ev A.N.†

*Department of Theoretical Physics, St.-Petersburg University,
Ul'yanovskaya 1, Staryi Petergof, 198904, St.-Petersburg, Russian Federation*

Abstract

Using knowledge of the explicit n dependence of the RG functions and expressions of critical exponents in the framework of large N expansion in the Gross Neveu model we derive RG functions in 4- and 5-loop approximation.

*E-mail address: stepan at amoco.saclay.cea.fr

†This work was supported by the Grant of the American Physical Society

‡E-mail address: ioffe at onti.phys.lgu.spb.su

1 Introduction.

It is well known that the structure of renormalized field theory is determined by the renormalization group (RG) equations. One considerable problem then is the calculation of renormalization group functions. Usually one can compute only a few orders in perturbation theory. Hence, we do not have a complete knowledge of these functions. Other methods, such as the large N expansion, usually allow us to compute critical exponents at leading order only because calculations become too complicated at higher orders.

To overcome these difficulties other methods have been developed. One such approach uses critical conformal invariance at the fixed point of renormalization group for calculation of critical exponents, which are values of RG functions at fixed point. With this method it is possible to derive the critical exponents at order $1/N^3$. Using then the exact N dependence of RG functions at each order in perturbation theory one can reconstruct them at some high-loop approximation.

In this paper we derive five loop approximation for RG functions (up to few unknown coefficients) in the Gross Neveu and the non-linear σ models.

2 Preliminaries.

The $U(N)$ -symmetric massive Gross Neveu (GN) model describes a system of N D -dimensional Dirac spinors, its non-renormalized action being

$$S = \int d^D x [\bar{\psi}_a (\hat{\partial} + m_0) \psi_a + g_0 (\bar{\psi}_a \psi_a)^2 / 2] \quad (1)$$

$a = 1, \dots, N$ and summation over repeated indices is implied (in what follows we consider the theory in Euclidean space and functional distribution is denoted by $\exp S$ without the minus in the exponent). It is believed to be multiplicatively renormalizable in the framework of $2 + \varepsilon$ expansion and its renormalized action in dimension $D = 2 + 2\varepsilon$ is

$$S_R = \int d^D x [Z_0 m^2 M^{2\varepsilon} + \bar{\psi}_a (Z_1 \hat{\partial} + Z_2 m) \psi_a + g M^{-2\varepsilon} Z_3 (\bar{\psi}_a \psi_a)^2 / 2] , \quad (2)$$

where g is a dimensionless renormalized coupling constant, m is a renormalized mass, M is a renormalization mass and $Z_{0,1,2,3}$ are renormalization constants. The counterterms of model (2) (without vacuum one) are generated by the standard multiplicative renormalization procedure

$$\psi \rightarrow Z_\psi \psi , \quad m_0 = m Z_m , \quad g_0 = g Z_g M^{-2\varepsilon} , \quad (3)$$

$$Z_1 = Z_\psi^2 , \quad Z_2 = Z_\psi^2 Z_m , \quad Z_3 = Z_\psi^4 Z_g . \quad (4)$$

RG functions γ_F (the anomalous dimensions of quantities F) and β function are defined by the relations

$$\gamma_F = \tilde{\mathcal{D}}_\mu \ln Z_F \quad (F = \psi, m, g) , \quad \beta = \tilde{\mathcal{D}}_\mu g = (2\varepsilon g - \gamma_g)g , \quad \tilde{\mathcal{D}}_\mu \equiv \mu \frac{d}{d\mu} \Big|_{g_0, m_0} . \quad (5)$$

The β function of model (2) has an UV fixed point $\beta(g_*) = 0$ with $g_* \sim \varepsilon$. Critical dimensions $\Delta_F = d_F + \gamma_F^*$ (canonical + anomalous) of various quantities F (fields and parameters) in the framework of $2 + \varepsilon$ expansion are defined by the relations

$$\Delta_F = d_F + \gamma_F^* \ (F = \psi, m) \ , \quad \Delta_\tau \equiv 1/\nu = -\beta'(g_*) \ , \quad (6)$$

where $\gamma_F^* \equiv \gamma_F(g_*)$, $\Delta_\tau \equiv 1/\nu$ is the critical dimension of “temperature” $\tau = g - g_*$ and d_F are the corresponding canonical dimensions: $d_\psi = D/2 - 1$, $d_m = 1$.

For practical calculations it is more convenient to use the following model:

$$S = \int d^D x \ [\bar{\psi}_a(\hat{\partial} + m_0)\psi_a - \sigma^2/2g_0 + \sigma\bar{\psi}_a\psi_a] \ . \quad (7)$$

The generalization of this model which is multiplicatively renormalizable in the framework of $2 + \varepsilon$ expansion is

$$\bar{S} = \int d^D x \ [\bar{\psi}_a(\hat{\partial} + \bar{m}_0)\psi_a - \sigma^2/2 + v_0\sigma\bar{\psi}_a\psi_a + g_{20}(\bar{\psi}_a\psi_a)^2/2 + h_0\sigma] \quad (8)$$

with two independent bare charges g_{20} and $g_{10} \equiv v_0^2$. Its renormalized form is

$$\begin{aligned} \bar{S}_R = \int d^D x \ [& \bar{Z}_0\bar{m}^2M^{2\varepsilon} + \bar{\psi}_a(\bar{Z}_1\hat{\partial} + \bar{Z}_2\bar{m})\psi_a - \bar{Z}_3\sigma^2/2 + \\ & + \bar{Z}_4v_1M^{-\varepsilon}\sigma\bar{\psi}_a\psi_a + \bar{Z}_5g_2M^{-2\varepsilon}(\bar{\psi}_a\psi_a)^2/2 + \bar{Z}_6\bar{m}M^\varepsilon\sigma + h\sigma] \ . \end{aligned} \quad (9)$$

If we integrate model (9) over field σ we obtain model (2) with parameters:

$$m = \bar{m} + hvM^{-\varepsilon} \ , \quad g = g_1 + g_2 \ , \quad g_1 \equiv v^2 \quad (10)$$

and additional constant $\text{tr} \ln([\hat{\partial} + m]/[\hat{\partial} + \bar{m}]) + (h^2/2) \int d^D x$ to the action. Comparing counterterms in the models (2) and (9) one can obtain the following relations for renormalization constants

$$\left. \begin{aligned} \bar{Z}_1 = Z_1 \ , \quad \bar{Z}_3^{-1} = 1 + 2g_1(Z_0 + n/8\pi\varepsilon) \ , \quad \bar{Z}_2 = \bar{Z}_4 = Z_2\bar{Z}_3 \ , \\ v\bar{Z}_6 = 1 - \bar{Z}_3 = 2g_1(\bar{Z}_0 + n/8\pi\varepsilon) \ , \quad g_1\bar{Z}_3^{-1}\bar{Z}_4^2 + g_2\bar{Z}_5 = g\bar{Z}_3 \end{aligned} \right\} \quad (11)$$

where $g_1 \equiv v^2$ and n is defined by (12). These relations allow to express 7 constants $\bar{Z}_i(g_1, g_2)$ of model (9) through 4 constants $Z_i(g = g_1 + g_2)$ of model (2). It is more convenient to calculate the renormalization constants in the framework of massless ($\bar{m} = 0, h = 0$) one-coupling model like (9) with interaction $\sim \bar{\psi}\psi\sigma$ and additional four-fermion counterterm which breaks the multiplicative renormalizability.

To generate a systematic $1/n$ expansion one usually uses the model (7) with a scalar field σ . As have been shown in [5] the massless version of (9) possesses critical conformal invariance at the D -dimensional fixed point of renormalization group and the exponent η have been calculated up to order $1/n^3$ with the conformal bootstrap method.

In what follows we shall consider $1/n$ expansions in arbitrary dimension of space D and the following notations will be used:

$$N \text{tr} \mathbf{1} \equiv n \ , \quad D \equiv 2\mu \ , \quad (12)$$

where $\text{tr } \mathbf{1}$ is a trace of the unit matrix in the space of D -dimensional spinors. In the framework of the $1/n$ expansion for the model (7) the canonical dimensions of the fields $\varphi \equiv \psi, \sigma$ and parameter τ are

$$d_\psi = d_{\overline{\psi}} = \mu - 1/2, \quad d_\sigma = 1, \quad d_\tau = 2(\mu - 1). \quad (13)$$

The notation $d[F] \equiv d_F$ means canonical dimension of quantity F . Let us parametrize the critical dimensions $\Delta_F = d_F + \gamma_F^*$ (canonical + anomalous) as follows:

$$\Delta_\psi = d_\psi + \eta/2, \quad \Delta_\sigma = d_\sigma - \eta - 2\Delta, \quad \Delta_\tau \equiv 1/\nu \equiv 2\lambda \quad (\lambda_0 = \mu - 1), \quad (14)$$

where η and ν are the usual notations for the critical exponents, the quantity 2Δ is a critical dimension of the vertex $\overline{\psi}\psi\sigma$ in the model (7). For this model the quantity Δ_τ is connected with the critical dimension $\Delta[\sigma^2]$ of a composite operator $F = \sigma^2$ by means of the following relation

$$\Delta_\tau \equiv 1/\nu \equiv 2\lambda = 2\mu - \Delta[\sigma^2]. \quad (15)$$

For the $1/n$ expansion of any exponent z its coefficients in powers of $1/n$ are denoted by z_k : $z = z_0 + z_1/n + z_2/n^2 + \dots$. The first coefficients $\eta_1, \Delta_1, \lambda_1$ for the special case of dimension $D = 2\mu = 3$ have been calculated in [1]. The coefficient η_1 for arbitrary dimension $D = 2\mu$ have been calculated in [2], Δ_1 in [3] and λ_1, η_2 in [4]. In [5] coefficients Δ_2, λ_2 and η_3 have been calculated.

The expression for η_3 contains function $I(\mu)$ which cannot be expressed explicitly through the gamma-function $\Gamma(z)$ and its derivatives. The function $I(\mu)$ is expressed through the self-energy massless diagram showed in Fig.1 on which a line with index a means a massless propagator $|x - y|^{-2a}$. According to its dimension the diagram is a simple line with index $\mu - 1 + \Delta$ multiplied by the coefficient $\Pi(\mu, \Delta)$ which is the “value” of the diagram. The quantity $I(\mu)$ is defined from $\Pi(\mu, \Delta)$ by the relation $I(\mu) = d \ln \Pi(\mu, \Delta) / d\Delta|_{\Delta=0}$ (for $\Delta = 0$ the diagram is calculated exactly: $\Pi(\mu, 0) = 3\pi^{2\mu}(\psi'(\mu - 1) - \psi'(1))\Gamma(\mu - 1)\Gamma(2 - \mu)/\Gamma(2\mu - 2)$). In [6] the value of $I(\mu)$ for the dimension $D = 2\mu = 3$ and the first terms of ϵ expansions around $D = 4$ are given: $I(3/2) = 3\psi''(1/2)/2\pi^2 + 2 \ln 2$, $I(2 - \epsilon) = O(\epsilon)$ and in [6, 7] the ϵ expansion around $D = 2$ was obtained

$$I(1 + \epsilon) = -\frac{2}{3\epsilon} + \frac{2\zeta(3)\epsilon^2}{3} - \zeta(4)\epsilon^3 + \frac{13\zeta(5)\epsilon^4}{3} + \dots \quad (16)$$

An alternative way of derivation of ϵ expansion for $I(\mu)$ is presented in the Appendix.

Using $1/n$ expansion of critical exponents and some additional information about the explicit n dependence of RG functions one can derive high loop approximation for RG functions. This additional information usually includes the knowledge of exact expressions of the RG functions for some special values of n .

3 Five loop approximation for the Gross Neveu and the σ models.

As have been noted in [8] the case $n = 1$ ($N = 1/2$) corresponds to Majorana spinor and, hence, there are no four-fermion interaction term in (1), (2), (8), (9) and we have

for model (2)

$$Z_0 = 0, Z_1 = Z_2 = 1 \quad (17)$$

and Z_3 is undefined as coefficient of zero.

Note that though the interaction $(\bar{\psi}\psi)^2$ does not exists at $n = 1$ but $\sigma\bar{\psi}\psi$ does and for (9) we have

$$\bar{Z}_1 = 1, \quad \bar{Z}_2^{-1} = \bar{Z}_3^{-1} = \bar{Z}_4^{-1} = 1 + \frac{\lambda^2}{4\pi\varepsilon}, \quad \bar{Z}_0 = -\frac{\lambda^2}{32\pi\varepsilon}\bar{Z}_2, \quad \bar{Z}_6 = \frac{\lambda}{4\pi\varepsilon}\bar{Z}_2 \quad (18)$$

but again we can say anything about the value of $\bar{Z}_5$ at $n = 1$.

From above considerations we conclude that for the model (2) $\gamma_\psi(g)|_{n=1} = \gamma_m(g)|_{n=1} = 0$ and, hence, these RG functions should contain the factor $(n - 1)$ at each order of perturbation theory. But we cannot say the same about the β function.

The case $n = 2$ corresponds to $N = 1, D = 2$. As have been noted very early the $U(1)$ Thirring and $N = 1$ Gross Neveu model are equivalent in the two dimensions. The Thirring model in the two dimensions is known to be exactly soluble and it is easy to derive the RG functions ($\varepsilon = 0, N = 1$):

$$\beta(u) = 0, \quad \gamma_\psi(u) = \frac{u^2}{1 + 2u}, \quad u \equiv \frac{g}{4\pi}. \quad (19)$$

But from [4] it is known the following three loop approximation for RG functions in the model (2):

$$\begin{aligned} \gamma_\psi &= u^2(n - 1) - u^3(n - 1)(n - 2) + O(u^4) \\ \beta(u) &= -2u^2(n - 2) + 4u^3(n - 2) + 2u^4(n - 2)(n - 7) + O(u^5) \end{aligned} \quad (20)$$

setting $n = 2$ in it we obtain

$$\beta|_{n=2} = 0, \quad \gamma_\psi|_{n=2} = u^2 + O(u^4). \quad (21)$$

From the above results we can say that in the framework of $2 + \varepsilon$ expansion the Thirring and the Gross Neveu models are not completely equivalent. Hence, there are no reason to believe that higher loop contributions of γ_ψ should contain the factor $n - 2$.

Using now this information and results of [5] it is possible to reconstruct RG functions of the Gross Neveu model in five loop approximation (up to some unknown coefficients) by expanding the critical exponents in both ε and $1/n$ and comparing then the corresponding coefficients. The result is the following:

$$\begin{aligned} \gamma_\psi(u) &= u^2(n - 1) - u^3(n - 1)(n - 2) + u^4(n - 1)[n^2 - 7n + 7] + \\ &+ u^5 \frac{(n - 1)}{3} [3n^3(-2\zeta(3) - 1) + 2n^2(-3\zeta(3) + 17) \\ &+ 12n(-11\zeta(3) - 10) + \eta_{50}] \end{aligned} \quad (22)$$

$$\begin{aligned}
\beta(u) = & 2u\varepsilon - 2u^2(n-2) + 4u^3(n-2) + 2u^4(n-2)(n-7) + \\
& + \frac{4}{3}u^5(n-2)[-n^2 - n(66\zeta(3) + 19) + \beta_{40}] + \\
& + \frac{1}{3}u^6(n-2)[3n^3(-2\zeta(3) + 1) + n^2(261\zeta(4) - 168\zeta(3) - 83) + \\
& + n\beta_{51} + \beta_{50}]
\end{aligned} \tag{23}$$

$$\begin{aligned}
\gamma_m(u) = & 2u(n-1) - 2u^2(n-1) - 2u^3(n-2)(2n-3) + \\
& + 2u^4 \frac{(n-1)}{3} [n^2(-6\zeta(3) + 5) + n(90\zeta(3) + 23) + m_{40}] + \\
& + u^5 \frac{(n-1)}{3} [3n^3(-6\zeta(3) - 5) + n^2(-81\zeta(4) + 264\zeta(3) + 169) + \\
& + nm_{51} + m_{50}]
\end{aligned} \tag{24}$$

Where

$$\gamma_m = -\gamma[\bar{\psi}\psi] = 2\varepsilon - \gamma_\sigma \tag{25}$$

is the anomalous dimension of the mass and $\eta_{50}, \beta_{40}, \beta_{51}, \beta_{50}, m_{40}, m_{51}, m_{50}$ are unknown coefficients. In [4] the expression $(n-2)(n-5) = n^2 - 7n + 10$ for the four loop contribution of γ_ψ have been received instead $n^2 - 7n + 7$. This result have been derived in [4] using expressions for η_1, η_2 and due to the supposition that multiplier $(n-2)$ should be extracted. We checked our result (7 instead 10) by direct calculation of 30 relevant four loop diagrams in the MS scheme. So the supposition of extracting of multiplier $n-2$ to all orders in γ_ψ RG function is wrong.

Using a similar method and results of [9] it is possible to derive the following expressions for RG functions up to five loops in the non-linear σ model:

$$\begin{aligned}
\gamma_\varphi = & \frac{u}{2} - u^2 \frac{(n-2)}{2} + u^3 \frac{(n-2)(2n-5)}{8} + \\
& + u^4 \frac{(n-2)}{24} [-3n^2 + 2n + 14 + 3(3-n)(n+5)\zeta(3)] + \\
& + u^5 \frac{(n-2)}{192} [6n^3(3\zeta(4) + 4\zeta(3) + 2) - n^2(54\zeta(3) + 5) \\
& - n(315\zeta(4) + 1764\zeta(3) + 307) + a]
\end{aligned} \tag{26}$$

$$\begin{aligned}
\beta = & 2\varepsilon u - u^2(n-2) - u^3(n-2) - u^4 \frac{(n-2)(n+2)}{4} - \\
& - u^5 \frac{(n-2)}{12} [-n^2 + 22n - 34 + 18(n-3)\zeta(3)] + \\
& + u^6 \frac{(n-2)}{96} [3n^3(2\zeta(3) - 1) + n^2(81\zeta(4) + 144\zeta(3) - 11) + bn + c]
\end{aligned} \tag{27}$$

where a, b, c are unknown coefficients.

Appendix

In this appendix we give the main steps of derivation of recurrent relation for function $I(\mu)$. This method allow to obtain the expansion (16) without the procedure of summing infinite series of the Gegenbauer polinomials [7]. The definition of $I(\mu)$ is

$$I(\mu) \equiv \left. \frac{d}{d\Delta} \ln \Pi(\mu, \Delta) \right|_{\Delta=0}, \quad (\text{A.1})$$

where (see Fig.1)

$$\int d^D x_3 d^D x_4 x_{13}^{-2} x_{14}^{-2} x_{34}^{-2(\mu-1+\Delta)} x_{23}^{-2(\mu-1)} x_{24}^{-2(\mu-1)} = \Pi(\mu, \Delta) \cdot x_{12}^{-2(\mu-1+\Delta)}. \quad (\text{A.2})$$

Using transformation “ $\leftarrow$ ” from [9] the function $\Pi(\mu, \Delta)$ can be transformed to

$$\Pi(\mu, \Delta) = c \cdot F_\mu(2\mu - 3 + \Delta), \quad (\text{A.3})$$

where

$$\int d^D x_3 d^D x_4 x_{13}^{-2} x_{14}^{-2} x_{34}^{-2\alpha} x_{23}^{-2} x_{24}^{-2} = F_\mu(\alpha) \cdot x_{12}^{-2(\alpha+4-2\mu)} \quad (\text{A.4})$$

and

$$c = \frac{\Gamma(\mu - 1 + \Delta)\Gamma(\mu - 1 - \Delta)}{\Gamma(\mu - 1)\Gamma(\mu - 1)\Gamma(1 + \Delta)\Gamma(1 - \Delta)} = 1 + O(\Delta^2). \quad (\text{A.5})$$

From (A.5) it is obvious that the coefficient c is not contributed to $I(\mu)$ and we have

$$I(\mu) \equiv \left. \frac{d}{d\Delta} \ln F_\mu(2\mu - 3 + \Delta) \right|_{\Delta=0}. \quad (\text{A.6})$$

Using the Gram determinant method [10] one can obtain the following relation with shifted dimension of the space

$$-F_\mu(\alpha) - F_\mu(\alpha + 1) + F_\mu^{(1)}(\alpha) = \frac{2(2\mu - 1)(\mu - 1)}{\pi^2} F_{\mu+1}(\alpha + 2), \quad (\text{A.7})$$

where $F_\mu^{(1)}(\alpha)$ is a sum of integrals which can be integrated using the chain rule. This sum is

$$F_\mu^{(1)}(\alpha) = -4\pi^D [\alpha^2 + \alpha(-3\mu + 5) + \mu^2 - 5\mu + 5] \times \frac{\Gamma(\mu - 1)\Gamma(\mu - 1)\Gamma(\mu - 2 - \alpha)\Gamma(\alpha + 3 - 2\mu)}{\Gamma(\alpha + 2)\Gamma(3\mu - 3 - \alpha)}. \quad (\text{A.8})$$

On other hand, combining two recurrent relations [9] for the master diagram one can derive the following relation for $F_\mu(\alpha)$:

$$F_\mu(\alpha + 1) = -\frac{(2\mu - 3 - \alpha)}{(\mu - 2 - \alpha)} F_\mu(\alpha) + F_\mu^{(2)}(\alpha), \quad (\text{A.9})$$

where $F_\mu^{(2)}(\alpha)$ is again a sum of easily integrated diagrams:

$$F_\mu^{(2)}(\alpha) = -\frac{2\pi^D(3\mu-5-2\alpha)}{(2\mu-3-\alpha)(\alpha+2-\mu)^2} \frac{\Gamma(\mu-1)\Gamma(\mu-1)\Gamma(\mu-1-\alpha)\Gamma(\alpha+4-2\mu)}{\Gamma(\alpha+1)\Gamma(3\mu-4-\alpha)} . \quad (\text{A.10})$$

From (A.7) and (A.9) we get

$$F_\mu(\alpha) = \frac{2(2\mu-1)(\mu-\alpha-2)}{\pi^2} F_{\mu+1}(\alpha+2) + R_\mu(\alpha) , \quad (\text{A.11})$$

$$\begin{aligned} R_\mu(\alpha) &= \frac{(\mu-2-\alpha)}{(\mu-1)} [F_\mu^{(2)}(\alpha) - F_\mu^{(1)}(\alpha)] \\ &= -2\pi^D(\alpha+\mu) \frac{\Gamma(\mu-1)\Gamma(\mu-1)\Gamma(\mu-2-\alpha)\Gamma(\alpha+4-2\mu)}{\Gamma(\alpha+2)\Gamma(3\mu-3-\alpha)} . \end{aligned} \quad (\text{A.12})$$

Substituting now $\alpha = 2\mu - 3 + \Delta$ and accounting that at $\Delta = 0$ (A.2) can be integrated exactly (see, for example, [9]):

$$\Pi(\mu, 0) = F_\mu(2\mu-3) = 3\pi^D \frac{\Gamma(\mu-1)\Gamma(2-\mu)}{\Gamma(2\mu-2)} (\psi'(\mu-1) - \psi'(1)) , \quad (\text{A.13})$$

we derive the following relation for $I(\mu)$

$$I(\mu) = \frac{1}{(\mu-1)^2 A} \left[B + \frac{1}{6(\mu-1)} \right] + \left[1 + \frac{1}{(\mu-1)^2 A} \right] \left[I(\mu+1) + \frac{1}{\mu-1} \right] , \quad (\text{A.14})$$

where

$$A = \psi'(1) - \psi'(\mu-1) , \quad B = \psi(2\mu-2) + \psi(2-\mu) - \psi(\mu-1) - \psi(1) . \quad (\text{A.15})$$

Due to the property $I(2+\varepsilon) = \mathcal{O}(\varepsilon)$ [9, 5] one can obtain the expansion of $I(1+\varepsilon)$ ($D = 2 + 2\varepsilon$) in ε up to order ε^3 :

$$I(1+\varepsilon) = -\frac{2}{3\varepsilon} + \frac{2\zeta(3)}{3}\varepsilon^2 - \zeta(4)\varepsilon^3 + \mathcal{O}(\varepsilon^4) . \quad (\text{A.16})$$

References

- [1] G.Gat, A.Kovner, B.Rosenstein, Phys.Lett., 240B, 158, 1990.
- [2] S. Hikami, T. Muta, Prog. Theor. Phys. **57** (1977), 785.
- [3] J.Zinn-Justin, Nucl.Phys., B367, 105, 1991.
- [4] J.A.Gracey, Int.J.Mod.Phys., A6, 395, 1991,
J.A.Gracey, Int.J.Mod.Phys., A6, 2755, 1991.
- [5] A.N.Vasil'ev, S.E.Derkachov, N.A.Kivel, A.S.Stepanenko, Teor. Mat. Fiz., v.92, N3, 486–498 (1992); v.94, N2, 179–193 (1993); preprint T93/016, Saclay.
A.N.Vasil'ev, A.S.Stepanenko, “ $1/n$ –expansion in the Gross–Neveu Model: Computation of $1/\nu$ Exponent at Order $1/n^2$ with Conformal Bootstrap Method”, submitted to publication to Teor. Mat. Fiz.
- [6] A.N.Vasil'ev, Yu.M.Pis'mak, Yu.R.Khonkonen, Teor. Mat. Fiz., **50**, 195 (1982).
- [7] W. Bernreuther, F. Wegner, Phys. Rev. Lett. **57** (1986), 1383.
- [8] C.Luperini, P.Rossi, Ann.Phys., 212, 371–401, 1991.
- [9] A.N.Vasil'ev, Yu.M.Pis'mak, Yu.R.Khonkonen, Teor. Mat. Fiz., **46**, 157 (1980).
A.N.Vasil'ev, Yu.M.Pis'mak, Yu.R.Khonkonen, Teor. Mat. Fiz., **47**, 291 (1981).
- [10] S.E.Derkachov, J.Honkonen, Yu.M.Pis'mak. J. Phys. A: Math. Gen. **23** (1990) 5563–5576.




Fig.1